

**Children 11 Years and Older Need These Immunizations to Attend School in Arizona
2010-2011 School Year**

Immunizations	1st Dose	2nd Dose	3rd Dose	4th Dose	5th Dose
<u>(DTaP/DTP) Diphtheria, Tetanus & Pertussis, (DT) Diphtheria & Tetanus, (Td) Tetanus & Diphtheria</u>	Required	Required	Required	Required if dose #1 was given before 12 months of age.	
An interval of at least 4 weeks is required between the first doses of DTaP/DTP/DT/Td . An interval of 6 months is required between the last two doses.					
<u>(Tdap) Tetanus, Diphtheria & Pertussis</u>	Required for 6 th , 7 th , and 8 th graders who are 11 years and older and have had 5 years pass since their last DTaP/DTP//Td dose.				
Students 11 years and older in <u>all</u> grades are required to receive one dose of Tdap if 10 years have passed since their last dose of DTaP/DTP/DT/Td, or if they are in the process of receiving their first 3 doses of tetanus/diphtheria vaccine. Students who have received 1 dose of Tdap will need a Td booster after 10 years.					
<u>(MV/MCV) Meningococcal Vaccine</u>	Required for 6 th , 7 th , and 8 th graders who are 11 years and older.				
A dose of MV/MCV is recommended at 11 years of age. Because Meningococcal vaccine is licensed for use in children 2 years and older, doses given between 2 and 11 years of age are valid and acceptable.					
<u>(IPV/OPV) Polio</u>	Required	Required	Required	Required if the first three doses were all given before age 4.	
An interval of at least 4 weeks is required between all Polio doses. Refer children for the final dose 6 months after the previous dose, per new recommendations. <u>Children who received 4 Polio doses spaced 4 weeks apart are not required to repeat any doses.</u>					
<u>(MMR) Measles, Mumps, & Rubella</u>	Required	Required			
MMR #1 is due at 12-15 months of age. MMR #2 may be given 1 month after MMR #1 .					
<u>(Hep B) Hepatitis B</u>	Required	Required	Required. Must be at least 24 weeks of age.	Required only if dose #3 was given before 24 weeks of age.	
Hep B #1 may be given at birth. Dose #2 is due 4-8 weeks after dose #1. Dose #3 is due 2-5 months after dose #2, at least 4 months after dose #1 and no sooner than 24 weeks of age. A 2-dose adolescent series may be given to children 11-15 years of age over a 4-6 month period.					
<u>Varicella* (Chicken Pox)</u>	Required*	Required* if dose #1 was given at age 13 years or older. Varicella #2 is recommended, not required for all others.			
*Students who have history of chicken pox disease are not required to receive Varicella vaccine. Parental recall of chicken pox disease is acceptable. If MMR and Varicella vaccine doses were not given on the same day, they must be given at least 28 days apart.					